

Limit on the fermion masses in technicolor models

A. Doff and A. A. Natale

Instituto de Física Teórica, UNESP, Rua Pamplona 145, 01405-900, São Paulo, SP, Brazil
(February 1, 2008)

Recently it has been pointed out that no limits can be put on the scale of fermion mass generation (M) in technicolor models, because the relation between the fermion masses (m_f) and M depends on the dimensionality of the interaction responsible for generating the fermion mass. Depending on this dimensionality it may happen that m_f does not depend on M at all. We show that exactly in this case m_f may reach its largest value, which is almost saturated by the top quark mass. We make few comments on the question of how large can be a dynamically generated fermion mass.

The mechanism that breaks the electroweak gauge symmetry, $SU(2)_L \times U(1)_Y$, down to the gauge symmetry of electromagnetism $U(1)_{em}$ is the only still obscure part of the standard model. It is known that up to the scale of 1 TeV some sign of this mechanism has to become manifest in future experiments. In the same way that an upper bound on the scale of the electroweak symmetry breaking has been put forward (the 1 TeV scale), it was thought that the scale of fermion mass generation had also an upper bound, and that bound would be at reach of the next generation of accelerators [1]. Recently it was shown that no upper bound can be put on the scale of fermion mass generation beyond that on the scale of electroweak symmetry breaking [2]. This result was obtained considering the scattering of same helicity fermions into a large number of longitudinal weak vector bosons in the final state, and it was also obtained in a more involved way in Ref. [3]. This result is important and at the same time is disappointing, because an upper bound on the scale of fermion mass generation would provide a target for future accelerators in order to understand the origin of fermion masses. The scale related to the origin of fermion masses cannot be bounded but the fermion mass itself is bounded. The bound on the fermion masses comes out from the upper limit on the Yukawa coupling ($\lambda_y \leq \sqrt{8\pi}$) [4]. In the standard scenario this is not very interesting because it also leads to a bound in the fermion masses of the order of 1 TeV. Therefore there is still space for a heavy new family (respecting the constraints provided by the high precision experiments). This problem becomes much more interesting in theories with dynamical symmetry breaking like technicolor theories, where, in principle, some of the free parameters of the standard model are calculable as long as we know the symmetries of the underlying theory that is responsible for the mass generation. Let us recall some of the arguments about the nonexistence of a bound on the scale for fermion mass generation in technicolor models [2]. In these models the fermion mass is given by

$$m_f \approx c \frac{\langle \bar{\psi}_{tc} \psi_{tc} \rangle}{M_{etc}^2}, \quad (1)$$

where c is a constant and $\langle \bar{\psi}_{tc} \psi_{tc} \rangle$ is the technifermion condensate. M_{etc} is the mass of the extended technicolor boson and is the mass scale that reproduces an effective Yukawa coupling. According to Ref. [1] M_{etc} should be bounded in the following way. If technicolor is a QCD-like model we can assume $\langle \bar{\psi}_{tc} \psi_{tc} \rangle \approx v^3$ [5], where $v \approx 246$ GeV, and assuming $c \simeq g_{etc}^2$ we obtain

$$M_{etc} \approx \left(g_{etc}^2 \frac{v^3}{m_f} \right)^{1/2}, \quad (2)$$

which gives the bound on the mass scale responsible for fermion mass generation. Nowadays it is known that composite operators like $\langle \bar{\psi}_{tc} \psi_{tc} \rangle$ (and the technifermion self energy) may have a large anomalous dimension ($\gamma_m \gtrsim 1$) in such a way that the fermion mass is given by

$$m_f \approx c \frac{\langle \bar{\psi}_{tc} \psi_{tc} \rangle}{M_{etc}^2} \left(\frac{M_{etc}^2}{v^2} \right)^{\gamma_m}, \quad (3)$$

from where we notice that for $\gamma_m = 1$ there is no relation at all between m_f and M_{etc} , indicating the inexistence of a bound on this last mass scale. The anomalous dimension $\gamma_m = 1$ can be obtained in the extreme limit of a walking technicolor dynamics [6], corresponding to a near critical extended technicolor interaction with increased importance of four-fermion operators, and, if the M_{etc} scale is raised, $\gamma_m = 1$ possibly only happens with a fine tuning of the theory. Let us still continue to discuss the case $\gamma_m = 1$. Exactly in this case we cannot establish a bound on M_{etc} , but note that it also imply that the maximum dynamical fermion mass is limited by

$$m_f \leq cv. \quad (4)$$

In this note we propose to discuss which is the maximum value admitted by Eq.(4) or by the dynamical fermion mass in general. We will compute the dynamical fermion mass described in Fig.(1), where the ordinary fermions (f) are connected to technifermions (T_f) through an extended technicolor gauge boson associated to some gauge group ($SU(N_{etc})$ with coupling $\alpha_{etc} = g_{etc}^2/4\pi$.)




FIG. 1. Diagram for ordinary fermion dynamical masses in technicolor models.

We perform the calculation of Fig.(1) using the following general expression for the techniquark self-energy [7]

$$\Sigma(p)_g = \mu \left(\frac{\mu^2}{p^2} \right)^\theta [1 + b g_{tc}^2(\mu^2) \ln(p^2/\mu^2)]^{-\gamma \cos(\theta\pi)} \quad (5)$$

where in the last equation we identified $\gamma = \gamma_{tc}$. The scale μ (or v) is related to the technicolor condensate $\langle \bar{\psi}_{tc} \psi_{tc} \rangle \equiv \mu^3$ and is ultimately fixed by the experimental value of the weak gauge boson masses. The advantage of using such expression is that it interpolates between the extreme possibilities for the technifermion self-energy, *i. e.* when $\theta = 1$ we have the soft self-energy giving by

$$\Sigma_s(p) = \frac{\mu^3}{p^2} [1 + b g_{tc}^2(\mu^2) \ln(p^2/\mu^2)]^\gamma, \quad (6)$$

which is the one obtained when the composite operator $\langle \bar{\psi}_{tc} \psi_{tc} \rangle$ has canonical dimension. When $\theta = 0$ operators of higher dimension may lead to the hard self-energy expression

$$\Sigma_h(p) = \mu [1 + b g_{tc}^2(\mu^2) \ln(p^2/\mu^2)]^{-\gamma}, \quad (7)$$

where γ must be larger than $1/2$ and the self-energy behaves like a bare mass [8]. Therefore no matter is the dimensionality of the operators responsible for the mass generation in technicolor theories the self-energy can always be described by Eq.(5). In the above equations g_{tc} is the technicolor coupling constant and $\gamma = 3C_{tc}/16\pi^2 b$, where $C_{tc} = \frac{1}{2}[C_2(R_1) + C_2(R_2) - C_2(R_{\bar{\psi}\psi})]$, with the quadratic Casimir operators $C_2(R_1)$ and $C_2(R_2)$ associated to the $R.H$ and $L.H$ fermionic representations of the technicolor group, and $C_2(R_{\bar{\psi}\psi})$ is related to the condensate representation. $b_{tc} = \frac{1}{16\pi^2} [11N - \frac{2}{3}n_f]$ is the g_{tc}^3 coefficient of the technicolor group β function. The complete equation for the dynamical fermion mass is

$$m_f = \frac{3C_{etc}\mu}{16\pi^4} \int d^4q \left(\frac{\mu^2}{q^2} \right)^\theta \frac{g_{etc}^2(q) [1 + b_{tc} g_{tc}^2 \ln(\frac{q^2}{\mu^2})]^{-\delta}}{(q^2 + M_{etc}^2)(q^2 + \mu^2)}, \quad (8)$$

where C_{etc} is the Casimir operator related to the etc fermionic representations, a factor μ remained in the fermion propagator as a natural infrared regulator and $\delta = \gamma \cos \theta\pi$, $g_{etc}^2(q)$ is assumed to be giving by

$$g_{etc}^2(q^2) \simeq \frac{g_{etc}^2(M_{etc}^2)}{(1 + b_{etc} g_{etc}^2(M_{etc}^2) \ln(\frac{q^2}{M_{etc}^2})}. \quad (9)$$

Note that in Eq.(8) we have two terms of the form $[1 + b_i g_i^2 \ln q^2]$ where the index i can be related to tc or etc . To obtain an analytical formula for the fermion mass we will consider the substitution $q^2 \rightarrow \frac{x M_{etc}^2}{\mu^2}$, and we will assume that $b_{etc} g_{etc}^2(M_{etc}) \approx b_{tc} g_{tc}^2(M_{etc})$, what will simplify considerably the calculation. Knowing that the etc group usually is larger than the tc one we computed numerically the error in this approximation for few examples found in the literature. The resulting expression for m_f will be overestimated by a factor $1.1 - 1.3$ and is giving by

$$m_f \simeq \frac{3C_{etc} g_{etc}^2(M_{etc}) \mu}{16\pi^2} \left(\frac{\mu^2}{M_{etc}^2} \right)^\theta \left[1 + b_{tc} g_{tc}^2 \ln \frac{M_{etc}^2}{\mu^2} \right]^{-\delta} I, \quad (10)$$

where

$$I = \frac{1}{\Gamma(\sigma)} \int_0^\infty d\sigma \sigma^{\epsilon-1} e^{-\sigma} \frac{1}{\theta + \alpha\sigma},$$

and $\epsilon = \delta + 1 = \gamma \cos \theta\pi + 1$, $\alpha = b_{tc} g_{tc}^2(M_{etc})$. To obtain Eq.(10) we made use of the following Mellin transform

$$\left[1 + \kappa \ln \frac{x}{\mu^2} \right]^{-\epsilon} = \frac{1}{\Gamma(\epsilon)} \int_0^\infty d\sigma e^{-\sigma} \left(\frac{x}{\mu^2} \right)^{-\sigma\kappa} \sigma^{\epsilon-1}. \quad (11)$$

Finally we obtain

$$m_f \simeq \frac{3C_{etc} g_{etc}^2(M_{etc}) \mu}{16\pi^2} \left(\frac{\mu^2}{M_{etc}^2} \right)^\theta F(\cos \theta\pi, \gamma, \alpha). \quad (12)$$

where

$$F(\cos \theta\pi, \gamma, \alpha) = \left[1 + b_{tc} g_{tc}^2 \ln \frac{M_{etc}^2}{\mu^2} \right]^{-\gamma \cos(\theta\pi)} \Gamma(-\gamma \cos(\theta\pi), \theta/\alpha) \exp\left(\frac{\theta}{\alpha}\right) \alpha^{-1-\gamma \cos(\theta\pi)} \theta^{\gamma \cos(\theta\pi)}.$$

Simple inspection of the above equation shows that (as long as $M_{etc} > \mu$) the largest value for the fermion mass happens for $\theta = 0$, and expanding Eq.(12) near this point we have

$$m_f \simeq \frac{3C_{etc} g_{etc}^2(M_{etc}) \mu}{16\pi^2} \left[1 + b_{tc} g_{tc}^2 \ln \frac{M_{etc}^2}{\mu^2} \right]^{-\gamma} \times \frac{1}{\gamma b_{tc} g_{tc}^2(M_{etc})} (1 + O(\theta) + \dots) \quad (13)$$

and for $\theta = 0$ we obtain

$$m_f \simeq \frac{C_{etc} g_{etc}^2(M_{etc}) \mu}{C_{tc} g_{tc}^2(\mu)} \left[1 + b_{tc} g_{tc}^2 \ln \frac{M_{etc}^2}{\mu^2} \right]^{-\gamma+1} \quad (14)$$

which gives the largest dynamical fermion mass that we can generate. Although this result is simple and quite intuitive we have not been able to find it stated anywhere.

Note that using the expression for the running coupling $g_{etc}^2(M_{etc})$ Eq.(14) can be written in the following form:

$$m_f \sim \frac{C_{etc}}{C_{tc}} \left(\frac{\alpha_{etc}}{\alpha_{tc}} \right)^{\gamma_{tc}} \mu \sim c v \quad (15)$$

where γ of the previous expressions is indicating γ_{tc} , and the factor c is now giving by $c = \frac{C_{etc}}{C_{tc}} \left(\frac{\alpha_{etc}}{\alpha_{tc}} \right)^{\gamma_{tc}}$. The possible values of c will determine the maximum value of the fermion mass. To find some limits on the dynamical fermion mass let us consider some possible ways to introduce the extended technicolor theory. We may, for example, consider that the *etc* theory may be a kind of grand unified theory (gut) based on the group $SU(k)$ containing technicolor and the standard Georgi-Glashow group [9], we than have

$$SU(k) \supset SU(k-5)_{tc} \otimes SU(5)_{gg},$$

where $SU(k-5)_{tc}$ is the *tc* group, and $SU(5)_{gg}$ is the gut of Ref. [9]. As *tc* is a strongly interacting theory it is natural to have $k \geq 7$. Therefore, associating the $SU(k)$ group to *etc*, we obtain the following ratio of Casimir operators

$$\frac{C_{etc}}{C_{tc}} = \frac{(k^2 - 1)(k - 5)}{k((k - 5)^2 - 1)}.$$

On the other hand we must also preserve asymptotic freedom, what imply $k \leq 11$ [5] [12], and the ratio $r_c = \frac{C_{etc}}{C_{tc}}$ will take values in the range $r_c = 1.7 - 4.5$. We still have to look at the ratio of coupling constants. As *tc* is a QCD like theory we can assume as usual that $\alpha_{tc} \sim 1$. The *etc* theory can be associated to a gut in this case. Actually, there is no reason at all (specially when the self-energy is the expression with $\theta = 0$) to expect a low value for M_{etc} and a natural one could be $M_{etc} = \Lambda_{gut} \sim 10^{16}$ Gev with $\alpha_{etc} \approx \alpha_{gut} \sim (40)^{-1}$. The coefficient $\gamma_{tc} = \frac{3C_{tc}}{16\pi^2 b_{tc}}$ must be larger than 0.5. and in fact if the *tc* group is $SU(2)_{tc}$ we have $\gamma_{tc} \sim 0.5$, for other (and larger) models this coefficient will be larger than 1/2. Therefore we roughly have

$$\left(\frac{\alpha_{etc}}{\alpha_{tc}} \right)^{\gamma_{tc}} \sim \left(\frac{1}{40} \right)^{\frac{1}{2}} \sim \frac{1}{6}.$$

Finally, considering all the estimates we obtain

$$m_f^{max} \sim O(0.3 - 0.8)v \sim O(75 - 200)Gev. \quad (16)$$

Note that this is a rough estimative and possibly is the best that we can do considering the present knowledge of strongly interacting theories. Our calculation is possibly overestimated and it should be divided by a factor 1.1 to 1.3 as we informed in the paragraph after Eq. (9). We also assumed an extreme case for the self-energy maximizing the fermion mass and it is not clear if a realistic model can reproduce exactly this behavior. Therefore

considering only the smaller factor (1.1) discussed above it seems that the maximum value of the dynamical mass is already saturated by the top quark mass. There is a possible way to circumvent this limit, *i.e.* we could build a model with a fermion more massive than the limit given by Eq.(16) where the mass comes from the contribution of several diagrams. In this case the fermion mass could be given by $m_f = nm_f^{max}$, where n is the number of diagrams contributing to the mass of one specific fermion. Models of this kind are similar to the ones of Ref. [11] (a $SU(9)_{gut} \otimes SU(3)_H$ theory, with a technicolor gut and a horizontal symmetry group), which is based on the model of Ref. [10] (a $SU(7)$ technicolor gut). In the table below we show the maximum fermion mass that we can obtain in such models. In the $SU(9)$ model we have two diagrams feeding up the heaviest fermion, even so it is difficult to obtain a mass larger than the limit of Eq.(16). Note also that this result is quite dependent on the model and the introduction of a horizontal symmetry is necessary for building a realistic model and to give several contributions to the fermion masses.

$SU(k)$	$r_c = C_{etc=gut}/C_{tc}$	γ_{tc}	n	m_f^{max}
$SU(7)$	4.5	0.50	1	$O(177) Gev$
$SU(9)$	2.4	0.65	2	$O(110) Gev$

We can also consider a different class of models where the *etc* group (G_{etc}) and the standard model (G_{sm}) obey the following [13]

$$G_{etc} \otimes G_{sm} = SU(N_{etc}) \otimes SU(3)_c \otimes SU(2)_L \otimes U(1)_y, \quad (17)$$

where $SU(N_{etc})$ must be large enough to accommodate technicolor. No realistic model has yet been found along this line, but let us consider a model based on the $SU(5)_{etc}$ group [14] [15], which contain $SU(2)_{tc}$ and one technifermion generation. To obtain the hierarchy among the 3 generations the model has the following symmetry breaking structure

$$\begin{aligned} & SU(5)_{etc} \otimes G_{sm} \\ & \downarrow \Lambda_1 \sim 1000 Tev \\ & SU(4)_{etc} \otimes G_{sm} \\ & \downarrow \Lambda_2 \sim 100 Tev \\ & SU(3)_{etc} \otimes G_{sm} \\ & \downarrow \Lambda_3 \sim 10 Tev \\ & SU(2)_{tc} \otimes G_{sm} \end{aligned} \quad (18)$$

We will not discuss the details of this model but just assume that the *tc* dynamics has the behavior of Eq.(14) and compute the mass of the heaviest family wich is given by

$$m_3 \sim \frac{C_3}{C_{tc}} \left(\frac{\alpha_3(\Lambda_3)}{\alpha_{tc}} \right)^{\gamma_{tc}} v \sim c_3 v. \quad (19)$$

With the values discussed in Ref. [14] [15] we see that we do not obtain a very large mass and the limit of Eq.(16)

seems to be common to all models. It is also interesting to discuss another kind of constraint that can be put on the dynamical fermion masses. Fermion masses have been limited in the standard model making the analysis of the partial wave amplitudes ($J = 0$) of the processes $\bar{f}f \rightarrow \bar{f}f$ at high energies. These amplitudes will be proportional to $a_0 \propto m_f^2 G_F$, and the unitarity condition of the S matrix imply that $|a_0| \leq 1$, which gives the bound [16]

$$m_f^2 \lesssim \frac{2\pi\sqrt{2}}{3G_F}. \quad (20)$$

The question that we address now is if this limit can be directly applied to tc theories. If we follow Ref. [17] it seems that this result could be applied also to tc theories. In Ref. [17] it was shown that one technicolor theory where the dynamical symmetry breaking is generated due to the effect of higher order operators the resulting effective theory reproduces exactly the standard model (their self-energy solution is identical to the one we are discussing here). The gauge interactions of the ordinary fermions with the standard gauge boson is obviously the same, but more importantly the Higgs boson coupling is also reduced to the standard model one, what is fundamental to obtain the result of Eq.(20). Only a light degree of freedom (a scalar composite boson) appears below the TeV scale. Therefore the limit of Eq. (20) could be valid for technicolor when the tc dynamics is the harder one that is discussed here. Of course, this is not true for a softer self-energy solution, when the fermion mass is smaller due to the dependence on the scale M_{etc} . If we impose the limit of Eq.(20) over Eq.(15) we obtain

$$\frac{\alpha_{etc}(M_{etc})}{\alpha_{tc}(\mu)} \lesssim \left(\frac{2\pi\sqrt{2}}{3G_F} \frac{1}{v^2 r_c^2} \right)^{\frac{1}{2\gamma_{tc}}}. \quad (21)$$

Considering the numeric values $G_F \sim 1.166 \times 10^{-5} GeV^{-2}$, $v \sim 246 GeV$ and the ratio $r_c = 1.7 - 4.5$ we obtain $\alpha_{etc}(M_{etc})/\alpha_{tc}(\mu) < 1$. This limit will not imply in a very strong constraint in the dynamical fermion mass. However there is a problem in the above argument. If the dynamical symmetry breaking is generated by the effect of higher order operators the effective low energy theory may reproduce exactly the standard model as claimed in Ref. [17], but the effective Yukawa coupling will be also proportional to a form factor which, just on dimensional grounds, should be of the form $F(q^2 \rightarrow 0) \propto (1 - q^2/\mu^2 + \dots)$, because any other mass scale (like M_{etc}) is erased from the self-energy (or appears only in a logarithm). Therefore, at low energies the Yukawa coupling is equal to the one of the standard model, but for momenta q^2 near the tc scale (μ) this coupling should be quite suppressed leading to a dynamical fermion mass not higher than μ .

In conclusion, it seems very difficult to generate dynamical fermion masses in technicolor models larger than

the technicolor scale. The largest mass that can be obtained appears when we consider the hardest (concerning the momentum dependence) expression for the technicolor dynamics, which is the same that is also consistent with the nonexistence of a bound on the scale of fermion mass generation. Maybe models with some extra symmetry (possibly a horizontal symmetry), implying that the heaviest fermion receives mass contribution from several diagrams, could be one possibility to have fermions heavier than the top quark within the technicolor scheme, although we do not know any realistic model along this line. Otherwise, if technicolor is responsible for the standard model symmetry breaking, it seems that no other ordinary heavier fermion family will be found in the next generation of accelerators.

ACKNOWLEDGMENTS

This research was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (AAN) and by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) (AD).

-
- [1] T. Appelquist and M. S. Chanowitz, Phys. Rev. Lett. **59**, 2405 (1987).
 - [2] F. Maltoni, J. M. Niczyporuk and S. Willenbrock, Phys. Rev. **D65**, 033004 (2002).
 - [3] J. M. Cornwall, D. N. Levin and G. Tiktopoulos, Phys. Rev. Lett. **30**, 1268 (1973); Phys. Rev. **D10**, 1145 (1974).
 - [4] M. S. Chanowitz, M. A. Furman and I. Hinchliffe, Nucl. Phys. **B153**, 402 (1979); M. B. Einhorn and G. J. Goldberg, Phys. Rev. Lett. **57**, 2115 (1986); I. Lee, J. Shigemitsu and R. E. Shrock, Nucl. Phys. **B330**, 225 (1990).
 - [5] H. Georgi, Nucl. Phys. **B156**, 126 (1979).
 - [6] B. Holdom, Phys. Rev. **D24**, 1441 (1981); K. Yamawaki, M. Bando and K. I. Matumoto, Phys. Rev. Lett. **56**, 1335 (1986); M. Bando, K. I. Matumoto and K. Yamawaki, Phys. Lett. **B178**, 308 (1986); T. Appelquist, D. Karabali and L. C. R. Wijewardhana, Phys. Rev. Lett. **57**, 957 (1986); T. Appelquist and L. C. R. Wijewardhana, Phys. Rev. **D35**, 774 (1987).
 - [7] A. Doff and A. A. Natale, Phys. Lett. **B537**, 275 (2002).
 - [8] K. Lane, Phys. Rev. **D10**, 2605 (1974).
 - [9] H. Georgi and S. L. Glashow, Phys. Rev. Lett. **32**, 438 (1974).
 - [10] E. Farhi and L. Susskind, Phys. Rev. **D20**, 3404 (1979).
 - [11] A. Doff and A. A. Natale, hep-ph /0302166.
 - [12] M. T. Vaughn, J. Phys. **G5**, 1317 (1979).
 - [13] S. Dimopoulos and L. Susskind, Nucl. Phys. **B155**, 237 (1979).
 - [14] Thomas Appelquist and John Terning, Phys. Rev. **D50**, 2116 (1994).

- [15] Thomas Appelquist and Robert Shrock, Phys. Lett. **B548**, 204 (2002).
- [16] M. S. Chanowitz, M. A. Furman and I. Hinchliffe, Nucl. Phys. **B153**, 402 (1979); W. Marciano, G. Valencia and S. Willenbrock, Phys. Rev. **D40**, 1725 (1989).
- [17] J. Carpenter, R. Norton, S. Siegemund-Broka and A. Soni, Phys. Rev. Lett. **65**, 153 (1990).